

Universal Jump in the Helicity Modulus of the Two-Dimensional Quantum XY Model

Kenji Harada^a and Naoki Kawashima^b

^a *Division of Applied Systems Science, Kyoto University, Sakyo-ku, Kyoto 606-01, Japan*

^b *Department of Physics, Toho University, Miyama 2-2-1, Funabashi 274, Japan*

(11 February 1997)

The helicity modulus of the $S = 1/2$ XY model is precisely estimated through a world line quantum Monte Carlo method enhanced by a cluster update algorithm. The obtained estimates for various system sizes and temperatures are well fitted by a scaling form with L replaced by $\log(L/L_0)$, which is inferred from the solution of the Kosterlitz renormalization group equation. The validity of the Kosterlitz-Thouless theory for this model is confirmed.

The nature of phase transition of the quantum XY model in two dimensions has not been fully clarified while the existence of a transition at a finite temperature were suggested^{1,2} some years ago. Ding and Makivić³ examined this problem systematically by a large scale Monte Carlo simulation for the first time and concluded that a phase transition of the Kosterlitz-Thouless (KT)⁴ type takes place at $T_{KT} = 0.350(4)$ ³ or $0.353(3)$ ⁵. In their calculation, the temperature dependence of the correlation length and the linear susceptibility were studied. However, it is technically difficult to distinguish an exponential divergence from an algebraic one. Because of this difficulty, the validity of their conclusion on the nature of the phase transition was questioned⁶. For the same reason, Gupta and Baillie⁷ did not conclude, in spite of their very extensive Monte Carlo calculation, that the phase transition of the classical XY model is exactly what the KT theory predicts.

In previous studies, to our knowledge, a systematic study of the system size dependence of various quantities was missing for the quantum XY model. On the other hand, for other models with a KT phase transition, there are a number of reports on this size dependence. For example, Solýom and Ziman⁸ studied size dependence of the first excitation gap in the $S = 1/2$ anisotropic XXZ model in one dimension, which is exactly solvable and known to have a transition of the KT type at the anti-ferromagnetic isotropic point. They found that the exact estimates for the finite systems do not fit into the standard form of the finite-size scaling at the critical point. Instead of using the ordinary finite-size scaling form, Weber and Minnhagen⁹ used the Kosterlitz renormalization equation¹⁰ for the data analysis in their study of the classical XY model in two dimensions. They verified the KT type phase transition by comparing the size dependence of the helicity modulus, Υ , with the solutions of the renormalization group equation at the critical temperature. They found a remarkable agreement extending even to the prefactor of the logarithmic correction. Following the same idea, Olsson¹¹ performed a more detailed analysis of the classical model with a more extensive Monte Carlo simulation. He found the resulting data including off-critical ones consistent with the Kosterlitz renormalization group equation.

The helicity modulus is known to exhibit the universal jump at the critical temperature¹². This quantity corresponds to the super-fluid density when the model is regarded as a Boson system with hard-cores. In the world-line quantum Monte Carlo method¹³, the helicity modulus is related to the fluctuation in the total winding number of world-lines by the following equation¹⁴.

$$\Upsilon = (T/2)\langle \mathbf{W}^2 \rangle \quad (1)$$

where $\mathbf{W} \equiv (W_x, W_y)$ with W_x (W_y) being the total winding number in the x (y) direction. Makivić¹⁵ computed this quantity by means of a conventional world-line quantum Monte Carlo method. The method is not ergodic¹⁶, however, in that it does not allow the winding number to vary. Therefore, Makivić resorted to an alternative definition of the quantity although it does not results in exactly the same answer as the conventional definition does. Another difficulty of the conventional Monte Carlo method is its long auto-correlation time near and below the critical temperature. These difficulties limited the accuracy and the precision of the Makivić's estimates and narrowed the temperature-range of the simulation. To overcome these difficulties, we use the cluster Monte Carlo method¹⁶ in the present note, in which both the number of particles and the winding number can vary. Another advantage of the method is its autocorrelation time which is often shorter than that of the conventional method by several orders of magnitude.

In the present note, we attempt at a detailed and precise comparison between the model and the theory by Kosterlitz¹⁰ through an accurate estimation of the thermal fluctuation in the winding number near and below the critical temperature. We will see that such an estimation allows us to examine a new scaling form which is different from the ordinary finite-size scaling. In this scaling form, the distance from the critical point, i.e., $K - K_{KT}$ appears in the form of $(K - K_{KT})(\log(L/L_0))^2$, in contrast to $(T - T_{KT})L^{y_T}$ in the ordinary finite-size scaling. At the same time, the quantity $x \equiv \langle (\pi/2)\mathbf{W}^2 \rangle - 2$ scales as $x \log(L/L_0)$.

The $S = 1/2$ quantum XY model is defined by the following Hamiltonian.

$$H = -J \sum_{\langle ij \rangle} (S_i^x S_j^x + S_i^y S_j^y), \quad (2)$$

where $\langle ij \rangle$ runs over all nearest-neighbor pairs on the square lattice. As for the spin operators, we here use the convention in which $(S_i^\mu)^2 = 1/4$ ($\mu = x, y, z$).

In comparing the numerical result with the renormalization group theory, the squared winding number is a useful quantity since it can be regarded as the renormalized coupling constant and therefore its size dependence can be directly predicted by the following renormalization equation¹⁰.

$$\frac{dx}{dl} = -y^2, \quad \frac{dy}{dl} = -xy \quad (3)$$

Here, x and y are renormalized parameters after a renormalization operation up to the length scale $L \equiv L_0(T)e^l$ where $L_0(T)$ is some characteristic length of the order of the lattice constant and has no singularity at $T = T_{\text{KT}}$. The parameter x is related to the renormalized coupling constant, i.e., the helicity modulus by the following equation¹².

$$x = \frac{\pi \Upsilon}{T} - 2 = \frac{\pi}{2} \langle \mathbf{W}^2 \rangle - 2. \quad (4)$$

In what follows, we regard the estimate of $(\pi/2) \langle \mathbf{W}^2 \rangle - 2$ for a system of the size L at the temperature T as $x(l = \log(L/L_0(T)), T)$.

The equation (3) has an integral

$$\Delta(T) \equiv x^2(l) - y^2(l) \quad (5)$$

which does not depend on $l \equiv \log(L/L_0(T))$. As a function of T , this integral is non-singular even at $T = T_{\text{KT}}$, i.e., $\Delta(K) = a(K - K_{\text{KT}}) + b(K - K_{\text{KT}})^2 + \dots$ where $K \equiv J/T$. The solution of the renormalization equation (3) is given by

$$x(T, L) = \begin{cases} \sqrt{|\Delta|} \coth(\sqrt{|\Delta|}l) & (K > K_{\text{KT}}) \\ \sqrt{|\Delta|} \cot(\sqrt{|\Delta|}l) & (K < K_{\text{KT}}) \end{cases} \quad (6)$$

Note that the ordinary finite size-scaling form

$$x(T, L) = L^\omega g(\Delta L^{1/\nu}) \quad (7)$$

cannot be consistent with (6). Instead, the solution (6) is a special case of the following form that one can obtain from the ordinary finite-size scaling form by replacing L by $l = \log(L/L_0(T))$.

$$x(T, L) = l^{-1} f(\Delta l^2). \quad (8)$$

When expressed in the form of (8), the solution (6) corresponds to a ‘scaling’ function $f(X)$ which is non-singular at $X = 0$ in spite of the singularity in $\sqrt{|\Delta|}$ at $T = T_{\text{KT}}$. To be more specific, the solution (6) corresponds to

$$f(X) = 1 + O(X). \quad (9)$$

Note that the fitting functions used by Olsson¹¹ (Equations (11a-c) with (16) and (18) in his paper) are consistent with this scaling form. Olsson’s fitting function corresponds to a temperature-independent $L_0(T)$ in our notation.

For our Monte Carlo simulation, using Suzuki-Trotter decomposition, we transformed the two dimensional quantum XY model into the (2+1)-dimensional Ising model with four spin plaquette interactions. The partition function is described as

$$Z = \sum_S \prod_p \omega(S_p), \quad (10)$$

where S is the set of states $S_{i,t}^z$ on the (2+1)-dimensional lattice, S_p is a state of spins in a plaquette p formed by four spins $S_{i,t}^z, S_{j,t}^z, S_{i,t+1}^z, S_{j,t+1}^z$, and $\omega(S_p)$ is the two-spin propagator defined below. We apply periodic boundary conditions in all directions to preserve the translational invariance and to satisfy the trace requirements.

By numbering four local states on a bond as $1 = (\uparrow\uparrow)$, $2 = (\uparrow\downarrow)$, $3 = (\downarrow\uparrow)$ and $4 = (\downarrow\downarrow)$, the two-spin propagator can be written explicitly as

$$\omega = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cosh(\frac{\Delta\tau}{2}) & \sinh(\frac{\Delta\tau}{2}) & 0 \\ 0 & \sinh(\frac{\Delta\tau}{2}) & \cosh(\frac{\Delta\tau}{2}) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad (11)$$

where $\Delta\tau = J/(mk_{\text{B}}T)$ with the Trotter number m , and the temperature T . In what follows, we will set $J = 1$ and $k_{\text{B}} = 1$ for simplicity.

We have carried out a Monte Carlo simulation using the loop algorithm¹⁶. In this algorithm, four spins in each plaquette are connected pair-wise. The pairs are chosen stochastically. The connected spins form loops which are units of flipping. A world-line is a path connecting sites with up-spins which may wind around the system with periodic boundary condition. The winding numbers W_x (or W_y) is defined as the sum of the numbers of such windings of all world-lines in the x (or y) direction. For example, W_x can be rewritten as

$$W_x = \frac{1}{L_x} \sum_p \alpha_x(S_p) \quad (12)$$

where L_x is the lattice size in the x direction. The symbol $\alpha_x(S_p)$ stands for the function which takes the value 1 (or -1) if a world line passes through the plaquette p in the positive (or negative) x -direction and the value 0, otherwise. The other winding number W_y is defined in the same manner.

In order to reduce the statistical error, we have used an improved estimator for the squared winding number of world lines. It simply equals to one forth of the sum of squared winding numbers of loops formed by a clustering procedure (i.e., a graph assignment procedure) in the cluster Monte Carlo method. A more detailed discussion will be given elsewhere¹⁷. In the present simulation,

we have found that the new estimator reduces errors by about twenty percent.

In our simulations, we have taken various temperatures between 0.22 and 0.60 and used lattices with $L = 8, 16, 32$ and 64 . As the Trotter number m , we have used $m = 8, 16$ and 32 . When the systematic error due to the Trotter discretization exceeds the statistical error, we have reduced the systematic error by the extrapolation to $m = \infty$ using three data for different Trotter numbers. The length of a typical run on $L = 64$ at a temperature is more than 2×10^5 Monte Carlo Sweeps(MCS). To make use of the vector processors, we have developed the efficient vectorized code for the cluster identification. The underlying idea for this vectorization was proposed by Mino¹⁸ and it is based on the ‘divide-and-conquer’ strategy. In his algorithm, one firstly divides the lattice into small sub-lattices and, at the same time, identifies clusters in each sub-lattices neglecting the connectivity outside of the cluster. One then deals with twice larger sub-lattices formed by a pair of previous sub-lattices. By repeating this procedure, one can eventually get clusters identified correctly in the whole lattice. Using this algorithm, we have achieved the efficiency of 1.5 million site updates per second on the Fujitsu VPP500.

Each run is divided into several bins. The length of a bin is taken so that bin averages are statistically independent from each other at least approximately. We have measured integrated autocorrelation times for $\mathbf{W}^2$ at low temperatures $T < 0.35J$ by the standard binning analysis for a run of 10^5 MCS with $L = 64$. They turn out to be smaller than 2 MCS in all cases. Since the smallest bin length used in the main calculation is 1500 MCS, the statistical independence among bins is assured. The statistical errors of various observables have been calculated from the standard deviation of the bin-average distribution.

In Fig. 1, the raw data for $\langle \mathbf{W}^2 \rangle \equiv \langle W_x^2 + W_y^2 \rangle$ are plotted. Figure 2 is its ‘scaling plot’ using the form of (8) assuming that $L_0(T)$ does not depend on the temperature, which was a reasonably good approximation judging from the Olsson’s observation¹¹ (see Fig.14) for the classical model. ($L_0(T)$ in the present note corresponds to ‘const’ in Eq.(18) in Olsson’s paper.) The enlarged view around the critical point is shown as the inset. The scaling plot Fig. 2 is a reasonably good one, considering the fact that we have only two adjustable parameters, K_{KT} and L_0 , while there are three such parameters in the ordinary finite-size scaling plot. The value of K_{KT} is determined so that the resulting plot gives the ‘best’ scaling plot. To quantify the ‘goodness’ of the scaling plot, a cost function $S(K_{KT}, L_0)$ is defined¹⁹, which is essentially the deviation from the locally linear fitting. We used the data in the range $0 < \langle \mathbf{W}^2 \rangle - 4/\pi < 0.25$, which roughly corresponds to the temperature range $0.330 < T < 0.375$. The best fitting is obtained with the transition temperature

$$T_{KT} = J/K_{KT} = 0.342(2)J \quad (13)$$




FIG. 1. The helicity modulus (or super fluid density) $\Upsilon = (T/2)\langle \mathbf{W}^2 \rangle$ as a function of temperature. The universal jump is expected at the point where $\Upsilon = 2T/\pi$. Error bars are drawn but most of them are so small that they cannot be recognized.

which is significantly lower than the Ding and Makivić’s estimates^{3,5}. We confirmed that other choices of the range result in the estimates of the critical temperature consistent with the value quoted above. The best scaling plot is shown in Fig. 2. It should be noted that the scaling function takes the value 1 at $K - K_{KT} = 0$ as is predicted by the Kosterlitz renormalization group equation (see (9)). Of course, an ordinary finite-size scaling does not predict the value of the scaling function at this particular point. Considering the fact that we have not used this information while making the scaling plot, the agreement between the scaling theory and the numerical result is hardly understandable unless we assume the validity of the KT theory. (Note that the criterion we have adopted in choosing the value of K_{KT} is simply that all points should fall onto a single curve.) Therefore, we consider that the present result is a strong confirmation of the validity of the Kosterlitz’s scaling theory in the quantum XY model in two dimensions.

We have also tried another analysis following Weber and Minnhagen⁹. At each temperature, we assumed the following system size dependence of the helicity modulus.

$$\frac{\pi\Upsilon}{2T} = \frac{\pi}{4}\langle \mathbf{W}^2 \rangle = A(T) \left(1 + \frac{1}{2 \log(L/L_0(T))} \right) \quad (14)$$

This fitting form is correct at the critical point with $A(T_{KT}) = 1$ ((8) and (9)). Since the number of data at each temperature is small, the critical temperature cannot be determined as the one at which the fitting is best. Instead, we plot the coefficient $A(T)$ as a function of temperature (Fig. 3). The critical temperature, then, is estimated as the point at which $A(T)$ takes on the value 1. Linear fitting of the data yields




FIG. 2. A rescaled plot of the winding number fluctuation. The inset is a enlarged view of the critical point.

$$T_{KT} = 0.3423(3)J, \quad (15)$$

which is consistent with (13). (It is natural that here we have a more precise value than (13), since we have assumed (14), a more specific fitting function than (8).)

To summarize, the loop algorithm have proven to be very efficient in studying the winding number particularly at low temperatures, and we have confirmed that the phase transition in the quantum $S = 1/2$ XY model is of the KT type avoiding the subtle and technically difficult comparison between an exponential divergence and an algebraic one. We have demonstrated that not only the magnitude of the jump but also very fine points in the Kosterlitz-Thouless-Nelson theory are realized in the present model.

The authors would like to thank K. Nomura for useful comments and discussions. The computations in the present work were performed on Fujitsu's VPP500 at Kyoto University Data Processing Center and partially at ISSP, the university of Tokyo.




FIG. 3. The helicity modulus divided by the magnitude of its universal jump. The dashed vertical line indicates the critical temperature at which the solid line crosses the dashed horizontal line ($A = 1$). The solid line is determined by a linear fitting.

C. F. Baillie and J. Apostolakis, Phys. Rev. Lett. **61**, 1996 (1988); R. Gupta and C. F. Baillie, Phys. Rev. B **45**, 2883 (1992).

⁸ J. Sólyom and T. A. L. Ziman, Phys. Rev. B **30**, 3980 (1984).

⁹ H. Weber and P. Minnhagen, Phys. Rev. B **37**, 5986 (1987).

¹⁰ J. M. Kosterlitz, J. Phys. C **7**, 1046 (1974).

¹¹ P. Olsson, Phys. Rev. B **52**, 4526 (1995).

¹² D. R. Nelson and J. M. Kosterlitz, Phys. Rev. Lett. **39**, 1201 (1977).

¹³ See for example, *Quantum Monte Carlo Methods in Equilibrium and Nonequilibrium Systems* ed. M. Suzuki (Springer 1987).

¹⁴ E. L. Pollock and D. M. Ceperley, Phys. Rev. B **36**, 8343 (1987).

¹⁵ M. S. Makivić, Phys. Rev. B **46**, 3167 (1992).

¹⁶ H. G. Evertz, M. Marcu and G. Lana, Phys. Rev. Lett. **70**, 875 (1993); N. Kawashima, J. Stat. Phys. **82**, 131 (1996), and references cited therein.

¹⁷ K. Harada and N. Kawashima, unpublished.

¹⁸ H. Mino, Comp. Phys. Comm. **66**, 25 (1991).

¹⁹ N. Kawashima and N. Ito, J. Phys. Soc. Jpn. **62**, 435 (1993).

¹ E. Loh, Jr., D. J. Scalapino and P. M. Grant, Phys. Rev. B **31**, 4712 (1985).

² T. Onogi, Doctor Thesis (University of Tokyo, 1986).

³ H.-Q. Ding and M. S. Makivić, Phys. Rev. B **42**, 6827 (1990); M. S. Makivić and H.-Q. Ding, *ibid.* **43**, 3562 (1991).

⁴ J. M. Kosterlitz and D. J. Thouless, J. Phys. C **6**, 1181 (1973).

⁵ H.-Q. Ding, Phys. Rev. B **45**, 230 (1992).

⁶ J. Sznajd, Phys. Rev. B **45**, 489 (1992); H.-Q. Ding and M. S. Makivić, *ibid.* **45**, 491 (1992).

⁷ R. Gupta, J. DeLapp, G. G. Batrouni, G. C. Fox,